



Toxicity Assessment of F-T Jet Fuel (JP-8/SPK)

LT Vish Mokashi, PhD, MSC, USN

Naval Health Research Center/ EHEL

2729 R St., Bldg. 837, Area B

Wright Patterson Air Force Base, OH 45433

937-904-9473

Vishwesh.mokashi@wpafb.af.mil

CDR Gail Chapman, PhD, MSC, USN

Officer-in-Charge EHEL

David R. Mattie, PhD, DABT

AFRL/RHPB

NIDA 2009



Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE MAY 2009		2. REPORT TYPE		3. DATES COVERED 00-00-2009 to 00-00-2009	
4. TITLE AND SUBTITLE Toxicity Assessment of F-T Jet Fuel (JP-8/SPK)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Health Research Center/ EHEL,2729 R St., Bldg. 837, Area B,Wright Patterson AFB,OH,45433				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Presented at the NDIA Environment, Energy Security & Sustainability (E2S2) Symposium & Exhibition held 4-7 May 2009 in Denver, CO.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 39	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

The views expressed are those of the presenter and do not reflect the official policy or position of the Department of the Navy, Department of Defense, or the U.S. Government. This presentation is approved for public release, distribution unlimited. This work was prepared as part of official duties. Title 17 U.S.C. §105 provides that “Copyright protection under this title is not available for any work of the United States Government.” Title 17 U.S.C. §101 defines a U.S. Government work as a work prepared by a military service member or employee of the U.S. Government as part of that person’s official duties.

Toxicity Assessment of F-T Jet Fuel (JP-8/SPK)

- Background on major jet fuels
- Background on JP-8/SPK
- Development of the JP-8/SPK toxicity program
- Dermal Irritation
- Genotoxicity
- Inhalation
- Conclusions
- Questions

Background

➤ DoD consumes ~ 5 B gallons of aviation jet fuel per year (US consumer ~25B)

➤ Major Jet Fuels:

- Jet-A
 - Commercial
- JP-4
 - Issued 1951
 - 50/50 kerosene-gasoline blend
 - Contains aliphatic and aromatic hydrocarbons, corrosion inhibitor and anti-static and -icing compounds
 - Phased out by USAF 1976-1996, replaced with JP-8
- JP-5
 - US Navy Fuel
 - Primarily kerosene
 - Lower volatility




Background

➤ Major Jet Fuels:

➤ JP-8 (MIL-DTL-83133)

- Kerosene-based replacement for JP-4
- Mixture of aliphatic, aromatic, and substituted naphthalene hydrocarbon compounds
- Reduced concentrations of benzene, n-hexane
- Contains icing inhibitor, corrosion inhibitors, lubricants, and anti-static agents
- +100 thermal stability additive can also contain metal deactivators, anti-oxidants



Erica Becvar
AFCEE/TDV

John Hinz, Health Risk
Assessment Support
USAF School of Aerospace
Medicine

Background



- Jet fuel toxicity risks traditionally associated with aromatic hydrocarbons
 - Conventional petroleum-based military aviation fuels contain ~20% Aromatics and 1-3% naphthalene.
- JP-4
 - toxicity studies have reported skin irritation, defatting, neurotoxicity nephrotoxicity and, in male rats, renal carcinogenicity.
 - In humans, symptoms of neurasthenia, psychasthenia, polyneuropathy and sexual dysfunction has been reported in military and civilian workers.

Background


➤ JP-8 Toxicity

- Considered slightly toxic
- Oral Lethal Dose (LD_{50}) > 5.0g/kg (highest tested)
- Dermal – non-irritating to slightly irritating
 - Repeated and continuous exposure does cause irritation in animals and humans
- Acute Vapor Inhalation Lethal Concentration (LC_{50}) > 3.43 mg/L (highest tested)
- Acute Aerosol LC_{50} > 4.39 mg/L (highest tested)
- Subchronic Inhalation- decreased body weight and renal hydrocarbon nephrothopy in male rats
- Chronic Inhalation – male rat nephrothapy



Background


- **JP-8/SPK (aka F-T, S-8...)**
 - USAF initiated the “alternative fuels” program in 1999
 - MIL-HDBK-510 Aerospace Fuels Certification
 - Goal of system certification and use of 50:50 synthetic jet fuel:JP-8 blend by 2016
 - To be used as a “drop in” fuel (JP-8:JP-8/SPK (50:50))
 - Synthetic jet fuel derived converting natural gas using coal-to-liquid (CTL) technology
 - Current synthetic jet fuels (S-8) alkane-rich, have reduced risk - essentially no aromatics or naphthalene




Background

➤ JP-8/SPK (aka F-T, S-8...)

- B-52 Certification (Aug 07)
- C-17 Ground/flight demos and FSE complete
- F101 engine (B-1B) test at AEDC
- CFM56 (F108) engine (KC-135) operability testing
- B-1B Demo flight – late Spr 08
- Acceleration of USAF Comm-Derivative aircraft certification by FAA Type-Cert (Dec 08?)
- F-22 certification effort - complete Fall 08
- F119 engine testing at P&W (May 08)
- F-16, KC-135, B-2 cert planning in-process



F-T Jet Fuel Toxicity Assessment



- Funded by Air Force
- Goal - Complete initial program by end FY08
- To ensure that toxicity program addresses all issues associated with testing a complex mixture, such as jet fuel, a review panel of toxicologists was established to discuss the toxicity testing program and provide advice on testing procedures.

F-T Jet Fuel Toxicity Assessment

➤ Panel members:

- David Mattie, Ph.D., DABT, Chair, AFRL/RHPB
- John Hinz, AFIOH/RSRE
- Gunda Reddy, PhD, USACHPPM
- LT Dean Wagner, PhD, MSC, USN, NHRC/EHEL
- David Steup, PhD, Shell Oil and Chairman of the Toxicology Committee, API
- Errol Zeiger, Ph.D., J.D., Errol Zeiger Consulting, Chapel Hill, NC

F-T Jet Fuel Toxicity Assessment

- **Develop appropriate study design for each toxicity test proposed:**
 - *In vitro* genotoxicity tests
 - Dermal irritation test
 - Acute inhalation study
 - *In vivo* genotoxicity test in tandem with inhalation rangefinder study
 - 90-day inhalation toxicity study

F-T Jet Fuel Toxicity Assessment

- All studies are completed.
- Final analysis and/or report preparation stage.
- Occupational exposure level will be developed by the review panel for the F-T fuel during the last quarter of FY09.

F-T Jet Fuel Toxicity Assessment

➤ Dermal Irritation Test

- Dermal Irritation Test
 - GLP Compliance
 - 100% concentrations of either SPK or JP-8
 - 50/50 mixture of SPK/JP-8
 - Occluded and semi-occluded exposures (each animal its own control)
 - Primary irritation index calculated to assign a descriptive rating to each test article and mixture

Dermal Toxicity

- ◆ JP-8 Occluded
- ◆ Primary Irritation Index = 2.1 Descriptive Rating = Moderately Irritating
- ◆ JP-8 Semi-Occluded
- ◆ Primary Irritation Index = 1.8 Descriptive Rating = Slightly Irritating
- ◆ SPK Occluded
- ◆ Primary Irritation Index = 2.3 Descriptive Rating = Moderately Irritating
- ◆ SPK Semi-Occluded
- ◆ Primary Irritation Index = 0.8 Descriptive Rating = Slightly Irritating
- ◆ JP-8/SPK Occluded
- ◆ Primary Irritation Index = 1.9 Descriptive Rating = Slightly Irritating
- ◆ JP-8/SPK Semi-occluded
- ◆ Primary Irritation Index = 1.5 Descriptive Rating = Slightly Irritating

F-T Jet Fuel Toxicity Assessment

In vitro genotoxicity tests

Performed by Eurofins/ Product Safety Labs

GLP Compliance

F-T Jet Fuel Toxicity Assessment

➤ Chromosomal Aberration (CA) Test

- Establish potential for fuel to induce chromosomal aberrations in human lymphocytes with and without activation.

➤ Conclusion

- Chromosomal aberrations were not induced by the fuel
- JP8/SPK is considered to be non-clastogenic

F-T Jet Fuel Toxicity Assessment

- **Reverse Mutation Assay (Ames Test)**
- Establish potential for fuel to induce gene mutations
- Detect point mutations in *Salmonella typhimurium*
 - Histidine reversion system His⁻ → His⁺

F-T Jet Fuel Toxicity Assessment

➤ Reverse Mutation Assay (Ames Test)

➤ Preliminary Results

- No toxic effects in any of five tester strains
- No relevant increase in revertants
- F-T jet fuel did not cause base pair or frame shift genetic mutations in any of the tester strains


➤ Conclusion

- F-T jet fuel is considered to be non–mutagenic in this bacterial reverse mutation assay


F-T Jet Fuel Toxicity Assessment

Inhalation studies

Protocol 07034: Generation of Jet Fuel: On/Off Valve in On Position



Protocol 07034: Generation of Jet Fuel: On/Off Valve in Off Position



F-T Jet Fuel Toxicity Assessment

➤ Acute Inhalation Study

➤ Completed:

- 4 hour exposure

- 2000 mg/m³ dose

- 5 males/5 females

- No clinical symptoms observed

F-T Fuel Toxicity

2-Week Study

- **2-week - Histological findings in:**
 - **lung: Inflammatory foci evident in highest doses**
 - **Olfactory epithelia: degeneration in highest doses**
 - **Kidney: hyaline droplet accumulation in all males exposed**
 - **Liver: Panlobular hepatocyte changes in all males exposed and females exposed to the highest concentration**

F-T Jet Fuel Toxicity Assessment

➤ 90-day Inhalation Study

➤ Study design based on:

- U.S. Environmental Protection Agency (U.S. EPA)
Harmonized Test Guideline developed by the Office of
Prevention, Pesticides and Toxic Substances (OPPTS)
870.3465 90-Day Inhalation Toxicity and
- Organization for Economic Cooperation and Development
(OECD) guideline 413, Subchronic Inhalation Toxicity:
- 90-day Study
- U.S. EPA Health Effects Test Guideline OPPTS 870.6200
Neurotoxicity Screening Battery.

F-T Jet Fuel Toxicity Assessment

➤ 90-day Inhalation Study

➤ Started April 28

➤ Route, Duration and Frequency of Administration

➤ Test substance will be administered as aerosol/vapor combination for 6 h/d, 5 d/w for 13 w

➤ Exposure Levels

➤ Animals will be exposed to three concentrations, high, intermediate and low, with a control group (0 mg/mm³).

➤ 2000 mg/m³, 700 mg/m³ and 200 mg/m³

F-T Jet Fuel Toxicity Assessment

➤ 90-day Inhalation Study

➤ Observations

- Animals will be observed before and after exposures for overt signs of toxicity
- Tissues and organs will be examined for gross pathology
- Blood will be collected for clinical pathology
- Tissues collected for histopathology including:
 - Nasal airways, trachea, larynx, lungs, liver, kidney, spleen, adrenals, heart and others

F-T Jet Fuel Toxicity Assessment

➤ 90-day Inhalation Study

- Motor Activity and Functional Observational Battery
 - To assess the neurobehavioral effects
- Sperm Morphology and Vaginal Cytology Examinations
 - Sperm will be stained and examined for percentage of abnormal sperm
 - Alterations in the estrous cycle will be assessed by examination cytology from vaginal smears

F-T Jet Fuel Toxicity Assessment

90-day Inhalation Study

Group	Exposure Level	Number of Animals	
		Males	Females
Control Replicate 1	0	5	5
Control Replicate 2		5	5
Low Replicate 1	200	5	5
Low Replicate 2		5	5
Intermediate Replicate 1	700	5	5
Intermediate Replicate 2		5	5
High Replicate 1	2000	5	5
High Replicate 2		5	5
Total		40	40

90-Day Inhalation Study

➤ Observations

- Animals observed before and after exposures for overt signs of toxicity
- Tissues and organs examined for gross pathology
- Blood collected for clinical pathology
- Full tissue list collected for histopathology
 - Analyzed male rat kidneys for alpha 2 microglobulin


90-Day Inhalation Study

Mean Exposure Levels


Group	Exposure Level	Number of Animals	
		Males	Females
Control	0.02 ± 0.10	10	10
Low	200.1 ± 5.0	10	10
Intermediate	698.6 ± 16.8	10	10
High	1988.4 ± 48.1	10	10

Body Weights

Protocol 08013 Females
DRAFT




Male Body Weights (without outliers)
DRAFT




Food Consumption

Protocol 08013 Average Daily Food Consumption, Females, mod
DRAFT



Protocol 08013 Average Daily Food Consumption, Males, mod
DRAFT



Hematology

- ◆ There were no biologically significant or toxicologically significant changes in hematology or clinical chemistry values seen in any of the groups.

Nasal Histopathology

Table 3 Incidence of selected findings in the noses of male rats (ten per group)

Group	Olfactory Epithelial Degeneration	Hypertrophy Hyperplasia Goblet Cells Nasopharyngeal Duct
Control	0	0
Low	0	0
Intermediate	9	10
High	10	10

Table 4 Incidence of selected findings in the noses of female rats (ten per group)

Group	Olfactory Epithelial Degeneration	Hypertrophy Hyperplasia Goblet Cells Nasopharyngeal Duct
Control	0	0
Low	0	0
Intermediate	8	9
High	10	10

Lung Histopathology

Table 5 Incidence of Inflammatory Cell Infiltration in the lungs of male rats (ten per group)

Group	Multifocal Inflammatory Cell Infiltration	Focal Inflammatory Cell Infiltration
Control	0	0
Low	0	1
Intermediate	0	1
High	10	0

Table 6 Incidence of Inflammatory Cell Infiltration in the lungs of female rats (ten per group)

Group	Multifocal Inflammatory Cell Infiltration	Focal Inflammatory Cell Infiltration
Control	0	0
Low	0	0
Intermediate	0	3
High	10	0

Kidney Histopathology

Table 7 Incidence of selected findings in the kidneys of male rats (ten per group)

Group	Increased Hyaline Droplet Tubular Epithelium
Control	0
Low	0
Intermediate	0
High	10

Alpha 2 Microglobulin

Group	Exposure Level	$\alpha_{2\mu}$ - globulin Concentration ($\mu\text{g}/\text{mg}$)	
		Males	Females
Control	0.02 ± 0.10	49.43 ± 20.87	0.16 ± 0.05
Low	200.1 ± 5.0	49.72 ± 20.57	N/A
Intermediate	698.6 ± 16.8	50.90 ± 13.72	N/A
High	1988.4 ± 48.1	35.65 ± 8.49	0.15 ± 0.01

F-T Jet Fuel Toxicity Assessment

Conclusions:

- **Dermal - animal protocol approved, start date TBD**
- **Genotox - No genetic mutations observed in vitro**
- **Inhalation**
 - **Acute - No clinical symptoms observed**
 - **2-week - Histological findings in:**
 - **lung: Inflammatory foci evident in highest doses**
 - **Olfactory epithelia: degeneration in highest doses**
 - **Kidney: hyaline droplet accumulation in all males exposed**
 - **Liver: Panlobular hepatocyte changes in all males exposed and females exposed to the highest concentration**
 - **90-day- Started 28 Apr 08**

F-T Jet Fuel Toxicity Assessment

Next:

- **Develop HHA**
- **Sensory Irritation Assay**
- **Genetic Biomarkers of Exposure**
- **Immunotoxicity Screen**

A silhouette of a fighter jet, likely an F-35, is positioned on the deck of an aircraft carrier. The scene is set at sunset or sunrise, with the sun low on the horizon, creating a bright, golden glow that reflects on the water. The jet's landing gear is visible, and it appears to be in the process of taking off or landing. The overall mood is dramatic and contemplative.

Questions?